
Bcl-10 Antibody

Subcategory: Mouse Monoclonal Antibody

Cat. No.: 251710

Unit: 0.1 mg

Description:

*****DISCONTINUED*****

B-cell lymphoma/leukemia 10 (Bcl-10) promotes apoptosis, pro-caspase-9 maturation and activation of NF-kappa-B via NIK and IKK. Bcl-10 is an adapter protein between the upstream TNFR1-TRADD-RIP complex and the downstream NIK-IKK-IKAP complex. Bcl-10 is a substrate for MALT1. Defects in Bcl-10 are involved in various types of cancer.

Isotype: Mouse IgG1

Applications: E, IHC, WB

Species Reactivity: H

Format: Each vial contains 0.1 mg IgG in 0.1 ml (1 mg/ml) of PBS pH7.4, 2% BSA with 0.09% sodium azide. Antibody was purified by Protein-G affinity chromatography.

Alternate Names: Bcl-10; B-cell lymphoma/leukemia 10; B-cell CLL/lymphoma 10; CED-3/ICH-1 prodomain homologous E10-like regulator; CIPER; CARD-containing molecule enhancing NF-kappa-B; Cellular homolog of vCARMEN; cCARMEN; Mammalian CARD-containing adapter molecule E10; mE10; Cellular-E10; c-E10; CARD-like apoptotic protein; hCLAP; BCL10; CIPER; CLAP

Accession No.: O95999

Antigen: Recombinant human Bcl-10.

Application Notes: E: 1:500-1:1,000; WB: 1:50-1:200; IHC: 1:25-1:100

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.