
Bfl-1 Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 253640

Unit: 0.1 mg

Description:

Apoptosis plays a major role in normal organism development, tissue homeostasis, and removal of damaged cells and is caused by caspase activation. Proteins that comprise the Bcl-2 family appear to control the activation of these enzymes. One such member is multi-domain antiapoptotic protein Bfl-1, which is overexpressed in stomach and other cancers. Bfl-1 can interact with Bax and suppress apoptosis by inhibiting the release of cytochrome c and caspase-3 activation. It is upregulated in cisplatin-resistant human bladder tumors, suggesting that its expression may be important for cisplatin resistance and inhibition of apoptosis in cancer cells. At least two isoforms of Bfl-1 are known to exist. Presumably due to post-translational modifications, Bfl-1 is often observed at higher molecular weight in SDS-PAGE than its predicted molecular weight would suggest.

Isotype: Rabbit Ig

Applications: E, WB, IHC

Species Reactivity: H, M

Format: Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.

Alternate Names: Bfl-1 ; Bcl-2-related protein A1; Hemopoietic-specific early response protein

Accession No.: NP_004040

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within human Bfl-1.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:00-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.