
Bif Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 253625

Unit: 0.1 mg

Description:

Apoptosis plays a major role in normal organism development, tissue homeostasis, and removal of damaged cells and is caused by the activation of proteolytic enzymes termed caspases. Proteins that comprise the Bcl-2 family such as Bax appear to control the activation of these enzymes. Bax activity was found to be regulated by its association with Bax-interacting factor 1 (BIF), a member of the endophilin B family that is associated with intracellular membranes. Following this interaction, Bax undergoes a conformational change and translocates to mitochondrial membranes. The Bax/BIF interaction appears to be enhanced by apoptotic stimuli, suggesting that BIF acts as the trigger to activate Bax, and as suppression of BIF promoted HeLa cell colony formation in soft agar, it may have a role in the suppression of cancer progression. At least two isoforms of BIF are known to exist.

Isotype: Rabbit Ig

Applications: E, WB, ICC

Species Reactivity: H, M

Format: Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.

Alternate Names: Bif; Bax-interacting factor 1; endophilin B1

Accession No.: AAK27365

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within human Bif.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; ICC: 1:100-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.