
CCR-7 Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 251667

Unit: 0.1 mg

Description:

C-C chemokine receptor type 7 (CCR-7) is a receptor for the MIP-3 beta chemokine. CCR-7 is a mediator of Epstein-Barr virus (EBV) effects on B-lymphocytes or of normal lymphocyte functions. CCR-7 is expressed in various lymphoid tissues and activated B- and T-lymphocytes. CCR-7 is strongly upregulated in B-cells infected with EBV and T-cells infected with herpesvirus 6 or 7.

Isotype: Rabbit Ig

Applications: E, WB, IHC

Species Reactivity: H, M, R

Format: Each vial contains 0.1 mg IgG in 0.1 ml (1 mg/ml) of PBS pH7.4 with 0.09% sodium azide. Antibody was purified by Protein-G affinity chromatography.

Alternate Names: CD197; CCR-7; C-C chemokine receptor type 7; C-C CKR-7; CC-CKR-7; MIP-3 beta receptor; Epstein-Barr virus-induced G-protein coupled receptor 1; EBF-induced G-protein coupled receptor 1; EBI1; BLR2; CDw197; CCR7; CMKBR7; EBI1; EVI1

Accession No.: P32248

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human CCR-7.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:100-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.