
CD3 (HIT3a) Antibody

Subcategory: Mouse Monoclonal Antibody

Cat. No.: 250934

Unit: 0.1 mg

Description:

The CD3 (HIT3a) antibody recognizes the 17-19 kD γ -chain of CD3 within the CD3 antigen/T cell antigen receptor (TCR) complex. The CD3 antigen is expressed in the cell cytoplasm during the early stage of T cell development and is expressed on the cell membrane at the late stage. CD3 antigen is displayed on 60-80% of normal peripheral blood lymphocytes and 60-70% of thymocytes and plays an important role in signal transduction after antigen recognition by TCR. The clone HIT3a has a strong mitogenic effect at ng level on T lymphocyte proliferation in soluble or immobilized conditions and has an immunosuppressive effect at high dose.

Conversely, the clone HIT3b has a strong mitogenic effect on T lymphocytes proliferation only under immobilized condition. NK cells also express the CD3 antigen in the cytoplasm, making the CD3 (HIT3a) antibody a helpful tool for analyzing the development relationship and the common precursor of both T cells and NK cells.

Isotype: mouse IgG2a

Applications: E, FC, IHC

Species Reactivity: H

Format: Each vial contains 0.1 mg IgG in 0.1 ml (1 mg/ml) of PBS pH7.4 with 0.09% sodium azide. Antibody was purified by Protein-G affinity chromatography.

Alternate Names: T-cell surface glycoprotein CD3; T-cell surface antigen T3/Leu-4; CD3; T3

Accession No.: P07766

Application Notes: Purified antibody is suitable for immunohistochemistry with acetone-fixed frozen sections. E: 1:500-1:1,000; FC: 1:200-1:1,000; IHC: 1:200-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

Product Citations: Shen DC. et al. 1993. ACTA Academia Medicinae Sinicae. 15(3):157; Schlossman S. et al., eds. 1995. Leucocyte Typing V: White Cell Differentiation Antigens. P246, Oxford University Press, New York.; Tadamitsu K. et al., eds. 1997. Leucocyte Typing VI: White Cell Differentiation Antigens. P49-52, 113-114 Garland Publishing, Inc., New York.

For research use only, not for diagnostic or therapeutic procedures.