

CstF-64 Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 200180

Unit: 0.1 mg

Description:

Cleavage stimulation factor 64 kDa subunit (CstF-64) is one of the multiple factors required for polyadenylation and 3'-end cleavage of mammalian pre-mRNAs. CstF-64 is directly involved in the binding to pre-mRNAs to form cleavage bodies, which are often adjacent to Cajal and PML bodies. The CSTF complex is composed of CSTF1 (50 kDa subunit), CSTF2 (64 kDa subunit) and CSTF3 (77 kDa subunit).

Isotype: Rabbit Ig

Applications: E, WB

Species Reactivity: H, M, X


Format: Each vial contains 0.1 mg IgG in 0.1 ml (1 mg/ml) of PBS pH7.4 with 0.09% sodium azide. Antibody was purified by Protein-G affinity chromatography.

Alternate Names: Cleavage stimulation factor 64 kDa subunit; CSTF 64 kDa subunit; CF-1 64 kDa subunit; CstF-64; CSTF2

Accession No.: P33240

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CstF-64.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500



The CstF-64 Antibody (Cat. No. 200180) is used in Western blot to detect CstF-64 in HEK293 cell lysate.

Storage: Store at -20°C. Minimize freeze-thaw cycles. Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.