
MD-2 (1A2E3) Antibody

Subcategory: Mouse Monoclonal Antibody

Cat. No.: 253271

Unit: 0.1 mg

Description:

MD-2 is a member of the Toll/interleukin-1 receptor (TIR) family, a group of proteins that include the Toll-like receptors (TLRs). TLRs are signaling molecules that recognize different pathogen-associated molecular patterns (PAMPs) and serve as an important link between the innate and adaptive immune responses. TLR4, the major signaling receptor for lipopolysaccharide (LPS), requires the binding of MD-2 to its extracellular region for maximal response to LPS. The specificity of this response is determined by the species of MD-2; e.g., human MD-2 transfected into mouse cells can cause mouse TLR4 to react to LPS analogs that are normally antagonistic to human but not mouse TLR4.

Isotype: mouse IgG1

Applications: E, WB, IHC

Species Reactivity: H, M, R

Format: Each vial contains 0.1 mg IgG in 0.1 ml (1 mg/ml) of PBS pH7.4 with 0.02% sodium azide. Antibody was purified by Protein-A affinity chromatography.

Alternate Names: MD-2; Lymphocyte antigen 96 precursor; ESOP-1

Accession No.: NP_056179

Antigen: Amino acids 2 to 160 of human MD-2.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:100-1:500

Storage: Store at 4°C. Minimize freeze-thaw cycles. Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.