
NADE Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 253485

Unit: 0.1 mg

Description:

The p75 neurotrophin receptor (p75NTR) is a member of the tumor necrosis receptor superfamily and can mediate cell death and cell survival in response to nerve growth factor (NGF). The p75NTR-associated cell death executor (NADE) mediates apoptosis by interacting with the cell death domain of p75NTR following the binding of NGF by p75NTR. Recent studies have shown that NADE also interacts with second mitochondria-derived activator of caspase (Smac). Co-expression of NADE and Smac promotes TRAIL-induced apoptosis and inhibits XIAP-mediated Smac ubiquitization. It has been suggested that it is this interaction between NADE and Smac that allows apoptosis to proceed. Finally, although initially discovered as an mRNA expressed in ovarian granulosa cells, NADE has been suggested to play a role in the neuronal death seen in epileptic brain damage.

Isotype: Rabbit Ig

Applications: E, WB, IHC

Species Reactivity: H, M

Format: Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by ion exchange chromatography.

Alternate Names: NADE

Accession No.: NP_996798

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within human NADE.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:00-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.