RICK Antibody

Subcategory: Rabbit Polyclonal Antibody
Cat. No.: 253351
Unit: 0.1 mg

Description:
Apoptosis is mediated by death domain (DD) and/or caspase recruitment domain (CARD) containing molecules and a caspase family of proteases. DD-containing serine/threonine kinase RIP regulates Fas-induced apoptosis. A novel CARD-containing serine/threonine kinase was recently identified and designated RICK/RIP2/CARDIAK for RIP-like interacting CLARP kinase, receptor interacting protein-2, and CARD-containing ICE associated kinase, respectively. RICK contains an N-terminal kinase catalytic domain and a C-terminal CARD domain. Overexpression of RICK induced apoptosis and activation of NF-kappaB and JNK. RICK interacts with members of the TRAF family, CLARP and caspase-1. Thus, RICK represents a novel kinase that regulates TNF and Fas induced-apoptosis and that is involved in the generation of proinflammatory cytokine IL-1beta. The messenger RNA of RICK is expressed in multiple human tissues.

Isotype: Rabbit Ig
Applications: E, WB, ICC
Species Reactivity: H
Format: Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.
Alternate Names: RICK ; RIP2
Accession No.: AF027706

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within human RICK.
Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; ICC: 1:100-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles. Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.