
TANK Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 253642

Unit: 0.1 mg

Description:

TANK was initially identified as a novel TRAF-interacting protein that regulated TRAF-mediated signal transduction. Specifically, ligand binding by surface receptors in the tumor necrosis factor (TNF) receptor and Toll/interleukin-1 (IL-1) receptor families lead to the formation of a TRAF/TANK complex that mediates the activation of the transcription factor NF-kappaB. This activation of NF-kappaB occurs through an association with the kinases IKKepsilon and TBK1. More recently, it was shown that these proteins can then form a complex with NEMO, a protein that regulates the activity of the IkappaB complex. This suggests that in addition to the possibility that TBK1 and IKKepsilon activate the IKKs, the association with the IKK complex may help these kinases modulate other functions, such as the transactivation potential of NF-kappaB proteins. At least two isoforms of TANK are known to exist.

Isotype: Rabbit Ig

Applications: E, WB, IHC

Species Reactivity: H, M, R

Format: Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.

Alternate Names: TANK ; TRAF family member-associated NF-kappaB activator; I-TRAF

Accession No.: NP_004171

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within human TANK.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:00-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.