TOCA-1 Antibody

Subcategory: Rabbit Polyclonal Antibody
Cat. No.: 253792
Unit: 0.1 mg

Description:
Actin reorganization is important for the regulation of neuronal morphology. A protein involved in this process, the transducer of cdc42-dependent actin assembly 1 (TOCA-1) protein, a member of the evolutionarily conserved pombe CDC15 homology (PCH) protein family, is an essential component of the Cdc42 pathway. TOCA-1 binds both N-WASP and Cdc42 and is essential for Cdc42- and PIP2-induced actin polymerization, suggesting that TOCA-1 mediates Cdc42-dependent actin nucleation by activating the N-WASP-WIP complex. Decreased expression of TOCA-1 significantly enhances neurite elongation in PC-12 cells; its overexpression in the same cells suppresses neurite elongation. It has been suggested that TOCA-1 negatively regulates axon branching by regulating membrane trafficking by regulating membrane trafficking through the F-BAR/EFC domain. Multiple isoforms of TOCA-1 are known to exist.

Isotype: Rabbit Ig
Applications: E, WB, IHC
Species Reactivity: H, M, R
Format: Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.
Alternate Names: TOCA-1 ; Transducer of cdc42-dependent actin assembly 1; formin binding protein 1-like; FNBP1L
Accession No.: NP_001020119

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within human TOCA-1.
Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:00-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles. Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.