

---

## **XEDAR Antibody**

**Subcategory:** Rabbit Polyclonal Antibody

**Cat. No.:** 253632

**Unit:** 0.1 mg

### **Description:**

X-linked ectodysplasin-A2 receptor (XEDAR) is a recently isolated member of the tumor necrosis factor receptor family that is highly expressed during embryonic development and binds to ectodysplatin-A2 (EDA-A2). Two predominantly expressed isoforms, XEDAR-s and XEDAR-L, differ by only a 21-amino region at the juxtamembrane region of the cytoplasmic domain. Neither isoform possesses a death domain and both have been shown to act mainly through TRAF3 and TRAF6 to activate the NF-kappaB and JNK pathways. Cells transfected with XEDAR and treated with EDA-A2 cause the assembly of a secondary complex containing FADD, caspase-8 and caspase-10, leading to the activation caspase-8 and caspase-3, and finally apoptosis. The EDA-A2-induced apoptosis is dependent on caspase-9 activation, as various pharmacological and genetic inhibitors of caspase-8 blocked apoptosis following EDA-A2 treatment.

**Isotype:** Rabbit Ig

**Applications:** E, WB, IHC

**Species Reactivity:** H

**Format:** Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.

**Alternate Names:** XEDAR; X-linked ectodysplasin-A2 receptor; EDA-A2 receptor; TNFRSF27

**Accession No.:** AAQ89953

**Antigen:** KLH-conjugated synthetic peptide encompassing a sequence within human XEDAR.

**Application Notes:** E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:00-1:500

**Storage:** Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.