

Cynomolgus CXCL12 / SDF-1 Protein (Fc Tag)

Catalog Number: 90009-C02H



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

CXCL12

Protein Construction:

A DNA sequence encoding the Cynomolgus (*Macaca fascicularis*) CXCL12 (NP_001036851.1) (Met 1-Lys 89) was fused with the Fc region of human IgG1 at the C-terminus.

Source: Cynomolgus

Expression Host: Human Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio-Activity

Measured by its ability to bind biotinylated recombinant human DPPIV in a functional ELISA.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Lys 23

Molecular Mass:

The recombinant cynomolgus CXCL12/Fc is a disulfide-linked homodimer. The reduced monomer consists of 309 amino acids and has a calculated molecular mass of 34.8 kDa. The apparent molecular mass of the reduced cynomolgus CXCL12/Fc monomer is approximately 39 kDa in SDS-PAGE under reducing conditions due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.


Usage Guide

Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

SDS-PAGE:



Reconstitution:

Detailed reconstitution instructions are sent along with the products.

Protein Description

Stromal cell-derived factor 1, also known as C-X-C motif chemokine 12, Interleukin 8, reduced in hepatomas, Pre-B cell growth-stimulating factor, SDF-1, PBSF, CXCL12 and SDF1, is a secreted protein which belongs to the interleukin alpha (chemokine CXC) family. CXCL12 / SDF-1 is widely expressed in various organs including heart, liver, brain, kidney, skeletal muscle, and lymphoid organs, and is highly conserved between species. CXCL12 / SDF-1 is expressed in two isoforms from a single gene that encodes two splice variants, SDF1α and SDF1β, which are identical except for the four residues present in the C-terminus of SDF1β but absent from SDF1α. As a highly efficacious lymphocyte chemoattractant, CXCL12 / SDF-1 activates and directs the migration of leukocytes in response to proinflammatory stimuli such as LPS, TNF, or interleukins. It has been demonstrated that CXCL12 / SDF-1 promotes tumor growth and malignancy, enhances tumor angiogenesis, participates in tumor metastasis, and thus plays an important role in carcinoma pathogenesis. In addition, CXCL12 / SDF-1 is identified as the biological ligand for CXCR4, which is widely expressed in leukocytes and acts as the cofactor for HIV-1 entry, and inhibits the HIV-1 infection. Furthermore, SDF1 is rapidly inactivated via N-terminal processing by cathepsin G associated with the membrane of B-cells, NK cells and to a lesser extent, T-cells.

References

1. Bleul, C.C. et al., 1996, Nature. 382: 829-833.
2. Sapede, D. et al., 2005, Proc. Natl. Acad. Sci. USA. 102: 1714-1718.
3. Delgado, M.B. et al., 2001, Eur. J. Immunol. 31: 699-707.
4. Orimo, A. et al., 2005, Cell. 121: 335-348.
5. Kryczek, I. et al., 2007, Am. J. Physiol. Cell. Physiol. 292: C987-995.
6. Bbalabanian, K. et al., 2005, J. Biol. Chem. 280: 35760-35766.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>