

Mouse Granulin / GRN / Progranulin Protein (His Tag)

Catalog Number: 50396-M08H



Sino Biological Inc.

Biological Solution Specialist

General Information

Gene Name Synonym:

RP23-226P4.4, epithelin

Protein Construction:

A DNA sequence encoding the mouse GRN (NP_032201.2) (Met 1-Leu 589) was fused with a polyhistidine tag at the C-terminus.

Source: Mouse

Expression Host: Human Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Thr 18

Molecular Mass:

The secreted recombinant mouse GRN comprises 583 amino acids and has a predicted molecular mass of 63 kDa. As a result of glycosylation, the apparent molecular mass of rmGRN is approximately 70-90 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.


Usage Guide

Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

SDS-PAGE:



Reconstitution:

Detailed reconstitution instructions are sent along with the products.

Protein Description

Granulin (GRN), also known as PC cell derived growth factor (PCDGF), is a ubiquitously expressed 88 kDa secreted, glycosylated protein which contains seven tandem repeats (55-57 aa consensus) of a highly conserved 12-cysteine granulin/epithelin motif. Cleavage of the signal peptide produces mature granulin which can be further cleaved into a variety of active, 6 kDa peptides. These smaller cleavage products are named granulin A, granulin B, granulin C, etc. Both the peptides and intact granulin protein regulate cell growth. Different members of the granulin protein family may act as inhibitors, stimulators, or have dual actions on cell growth, and thus perform important functions in normal development, wound healing, and tumorigenesis. Defects in GRN are the cause of ubiquitin-positive frontotemporal dementia (UP-FTD), an autosomal dominant neurodegenerative.

References

1. Bateman, A. et al., 1991, Biochem. Biophys. Res. Commun. 173 (3): 1161-8.
2. Hoque, M. et al., 2003, Mol. Cell. Biol. 23 (5): 1688-702.
3. Hoque, M. et al., 2005, J Biol Chem. 280 (14): 13648-57.
4. Zhou, Ying. et al., 2008, Cell. Signal. 20 (5): 942-8.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax : +86-10-51029969 • Tel: +86-400-890-9989 • <http://www.sinobiological.com>