

Human NRG1-alpha Protein (ECD, Fc Tag)

Catalog Number: 13499-H02H



Sino Biological Inc.

Biological Solution Specialist

General Information

Gene Name Synonym:

ARIA, GGF, GGF2, HGL, HRG, HRG1, HRGA, MST131, NDF, SMDF

Protein Construction:

A DNA sequence encoding the human NRG1 isoform alpha (Q02297-1) extracellular domain (Ser 20-Lys 241) was fused with the Fc region of human IgG1 at the C-terminus.

Source: Human

Expression Host: Human Cells

QC Testing

Purity: > 87 % as determined by SDS-PAGE

Bio-Activity

1. Measured by its ability to biotinylated Human ErbB4-Fc (cat:10363-H03H) in functional Elisa.
2. Measured by its ability to biotinylated Rhesus ErbB3-His (Cat:90043-K08H) in functional Elisa.
3. Measured by its ability to biotinylated Rhesus ErbB3 (Cat:90043-K02H) in functional Elisa.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Ser 20

Molecular Mass:

The recombinant human NRG1/Fc chimera is a disulfide-linked homodimeric protein. The reduced monomer consists of 463 amino acids and has a calculated molecular mass of 51 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rh NRG1/Fc monomer is approximately 60 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.


Usage Guide

Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

SDS-PAGE:



Reconstitution:

Detailed reconstitution instructions are sent along with the products.

Protein Description

Neuregulin 1 or NRG1 is one of four proteins in the neuregulin family that act on the EGFR family of receptors. This growth factor was originally identified as a 44-kD glycoprotein that interacts with the NEU / ERBB2 receptor tyrosine kinase to increase its phosphorylation on tyrosine residues. NRG1 is a trophic factor that has been implicated in neural development, neurotransmission, and synaptic plasticity. NRG1 has multiple isoforms that are generated by usage of different promoters and alternative splicing of a single gene. Neuregulin 1 (NRG1) is essential for the development and function of multiple organ systems, and its dysregulation has been linked to diseases such as cancer and schizophrenia. NRG1 is a schizophrenia candidate gene and plays an important role in brain development and neural function. Schizophrenia is a complex disorder, with etiology likely due to epistasis.

References

1. Nicodemus KK. et al., 2010, Arch Gen Psychiatry. 67 (10): 991-1001.
2. Tan W. et al., 2007, J Biol Chem. 282 (33): 24343-51.
3. Holmes WE. et al., 1992, Science. 256 (5060): 1205-10.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>