

Human SHH / Sonic hedgehog Protein (aa 1-197, His Tag)

Catalog Number: 10372-H08H1



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

HHG1, HLP3, HPE3, MCOPCB5, SMMCI, TPT, TPTPS

Protein Construction:

A DNA sequence encoding the amino acid sequence (Met 1-Gly 197) of human SHH (Q15465), that is Sonic hedgehog protein N-product, was fused with a polyhistidine tag at the C-terminus.

Source: Human

Expression Host: Human Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio-Activity

Measured by its ability to induce alkaline phosphatase production by C3H10T1/2 mouse embryonic fibroblast cells. Nakamura, T. et al. (1997) Biochem. Biophys. Res. Commun.237:465.
The ED50 for this effect is typically 2-10 µg/mL.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Cys 24

Molecular Mass:

The recombinant human SHH (aa 1-197) consists of 185 amino acids and has a predicted molecular mass of 21 kDa. The apparent molecular mass of the protein is approximately 24 kDa in SDS-PAGE under reducing conditions due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.


Usage Guide

Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

SDS-PAGE:



Reconstitution:

Detailed reconstitution instructions are sent along with the products.

Protein Description

Sonic HedgeHog, also known as sonic hedgehog protein, belongs to the hedgehog family and is expressed in fetal intestine, liver, lung, and kidney. It cannot be detected in adult tissues. Sonic HedgeHog is a protein that is vital in guiding the early embryo. It has been associated as the major inductive signal in patterning of the ventral neural tube, the anterior-posterior limb axis, and the ventral somites. Sonic HedgeHog binds to the patched receptor, which functions in association with smoothened, to activate the transcription of target genes. In the absence of sonic HedgeHog, patched receptor represses the constitutive signaling activity of smoothened. Sonic HedgeHog also regulates another factor, the gli oncogene. Sonic HedgeHog intercellular signal is essential for a various patterning events during development: signal produced by the notochord that induces ventral cell fate in the neural tube and somites, and the polarizing signal for patterning of the anterior-posterior axis of the developing limb bud. Sonic HedgeHog exhibits both floor plate- and motor neuron-inducing activity. Defects in sonic hedgehog can cause microphthalmia isolated with coloboma type 5, triphalangeal thumb-polysyndactyly syndrome and holoprosencephaly type.

References

1. Ericson J. et al., 1997, Cold Spring Harb Symp Quant Biol. 62: 451-66.
2. Marigo V. et al., 1996, Proc Natl Acad Sci. 93 (18): 9346-51.
3. Stone DM. et al., 1996, Nature. 384: 129-34.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>