

# Human CXCL12 / SDF1b Protein (Fc Tag)

Catalog Number: 10118-H01H



**Sino Biological Inc.**  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

CXCL12, PBSF, SCYB12, SDF-1a, SDF-1b, SDF1, SDF1A, SDF1B, TLSF-a, TLSF-b, TPAR1

### Protein Construction:

A DNA sequence encoding the N-terminally truncated human SDF1 $\beta$  (NP\_000600.1) (Lys 22-Met 93) was expressed with the fused Fc region of human IgG1 at the N-terminus.

**Source:** Human

**Expression Host:** Human Cells

## QC Testing

**Purity:** > 95 % as determined by SDS-PAGE

### Bio-Activity

Measured by its ability to bind biotinylated recombinant human DPPIV in a functional ELISA.

### Endotoxin:

< 1.0 EU per  $\mu$ g of the protein as determined by the LAL method

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

**Predicted N terminal:** Glu 20

### Molecular Mass:

The recombinant human SDF1/Fc is a disulfide-linked homodimeric protein. The reduced monomer consists of 309 amino acids and predicts a molecular mass of 35.2 kDa. As a result of glycosylation, rhSDF1/Fc monomer is approximately 43 kDa in SDS-PAGE under reducing conditions.

### Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.


## Usage Guide

### Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

## SDS-PAGE:



## Reconstitution:

Detailed reconstitution

with the products.

## Protein Description

Stromal cell-derived factor 1, also known as C-X-C motif chemokine 12, Interleukin 8, reduced in hepatomas, Pre-B cell growth-stimulating factor, SDF-1, PBSF, CXCL12 and SDF1, is a secreted protein which belongs to the intercrine alpha (chemokine Cx-C) family. CXCL12 / SDF-1 is widely expressed in various organs including heart, liver, brain, kidney, skeletal muscle, and lymphoid organs, and is highly conserved between species. CXCL12 / SDF-1 is expressed in two isoforms from a single gene that encodes two splice variants, SDF1 $\alpha$  and SDF1 $\beta$ , which are identical except for the four residues present in the C-terminus of SDF1 $\beta$  but absent from SDF1 $\alpha$ . As a highly efficacious lymphocyte chemoattractant, CXCL12 / SDF-1 activates and directs the migration of leukocytes in response to proinflammatory stimuli such as LPS, TNF, or interleukins. It has been demonstrated that CXCL12 / SDF-1 promotes tumor growth and malignancy, enhances tumor angiogenesis, participates in tumor metastasis, and thus plays an important role in carcinoma pathogenesis. In addition, CXCL12 / SDF-1 is identified as the biological ligand for CXCR4, which is widely expressed in leukocytes and acts as the cofactor for HIV-1 entry, and inhibits the HIV-1 infection. Furthermore, SDF1 is rapidly inactivated via N-terminal processing by cathepsin G associated with the membrane of B-cells, NK cells and to a lesser extent, T-cells.

## References

1. Bleul, C.C. et al., 1996, Nature. 382: 829-833.
2. Sapede, D. et al., 2005, Proc. Natl. Acad. Sci. USA. 102: 1714-1718.
3. Delgado, M.B. et al., 2001, Eur. J. Immunol. 31: 699-707.
4. Orimo, A. et al., 2005, Cell. 121: 335-348.
5. Kryczek, I. et al., 2007, Am. J. Physiol. Cell. Physiol. 292: C987-995.
6. Bbalabanian, K. et al., 2005, J. Biol. Chem. 280: 35760-35766.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>