

Cynomolgus ALK-2 / ACVR1 / ALK2 Protein (Fc Tag)

Catalog Number: 90058-C02H



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

ACVR1

Protein Construction:

A DNA sequence encoding the cynomolgus ACVR1 (F7A9J8) (Met1-Glu123) was expressed with the Fc region of human IgG1 at the C-terminus.

Source: Cynomolgus

Expression Host: Human Cells

QC Testing

Purity: (75.3+22.3) % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Asp 23

Molecular Mass:

The recombinant cynomolgus ACVR1 is a disulfide-linked homodimer. The reduced monomer comprises 342 amino acids and has a calculated molecular mass of 38.2 KDa. The apparent molecular mass of the protein is approximately 44 and 37 KDa respectively in SDS-PAGE.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.


Usage Guide

Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

SDS-PAGE:



Reconstitution:

Detailed reconstitution instructions are sent along with the products.

Protein Description

Cytokines and hormones belonging to the transforming growth factor beta (TGFβ) superfamily such as the activin, anti-müllerian hormone (AMH), bone morphogenetic proteins (BMPs), and Nodal, are involved in a variety of physiological processes including growth, cell differentiation, homeostasis, osteogenesis, and apoptosis. They initiate signaling through heteromeric complexes of a type I? and a type I I serine/threonine kinase receptor. The two type receptors are type I transmembrane proteins and are distinguished by the presence of a glycine/serine-rich juxta-membrane domain found in the type I receptors. Type II receptors are required for binding ligands and recruitment of type I receptors which then form stable complexes, and type I receptors are essential for signaling transduction. ALK2, also termed as ACVR1, ActRI, SKR1, was initially identified as an activin type I receptor because of its ability to bind activin in concert with ActRII or IIB. In addition, ALK2 is also identified as a BMP type I receptor. ACVR1 has been demonstrated to form complex with either the BMP-2/7-bound BMPR-II or ACVR2A/ACVR2B and transduce signals to downstream components R-SMADs. Recombinant soluble ALK2 does not bind activin. Mutations in this gene are associated with fibrodysplasia ossificans progressive.

References

1. Attisano L. et al., 1993, Cell. 75: 671-80.
2. Ten Dijke P. et al., 1994, Science. 264: 101-4.
3. Macias-Silva M. et al., 1998, J Biol Chem. 273: 25628-36.
4. Shore EM. et al., 2006, Nature Genetics. 38: 525-7.
5. Renlund N. et al., 2007, J Endocrinology. 195: 95-103.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>