

Human Apolipoprotein H / APOH Protein (His Tag)

Catalog Number: 11221-H08H



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

B2G1, BG, APOH

Protein Construction:

A DNA sequence encoding the human APOH (NP_000033.2) (Met 1-Cys 345) was expressed, with a polyhistidine tag at the C-terminus.

Source: Human

Expression Host: Human Cells

QC Testing

Purity: > 96 % as determined by SDS-PAGE

Bio-Activity

1. Measured by its binding ability in a functional ELISA.
2. Immobilized human APOH-his at 2 µg/mL (100 µl/well) can bind? biotinylated human LDLR-his, The EC50 of biotinylated human LDLR-his is 0.35 µg/mL.
3. Immobilized human APOH-his at 2 µg/mL (100 µl/well) can bind? biotinylated mouse LDLR-his, The EC50 of biotinylated mouse LDLR-his is 26 ng/mL.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gly 20

Molecular Mass:

The recombinant human APOH consists of 337 amino acids and predicts a molecular mass of 37.7 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rhAPOH is approximately 53 kDa due to different glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.


Usage Guide

Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

SDS-PAGE:



Reconstitution:

Detailed reconstitution instructions are sent along with the products.

Protein Description

Beta-2-glycoprotein 1, also known as Apolipoprotein H, Activated protein C-binding protein, APOH, B2GPI, and B2G1, is a glycoprotein synthesized by liver cells and it is present in the blood associated with plasma lipoproteins. APOH binds to various kinds of negatively charged substances such as heparin, phospholipids, and dextran sulfate. APOH may prevent activation of the intrinsic blood coagulation cascade by binding to phospholipids on the surface of damaged cells. APOH appears to completely inhibit serotonin release by the platelets and prevents subsequent waves of the ADP-induced aggregation. The activity of APOH appears to involve the binding of agglutinating, negatively charged compounds, and inhibits agglutination by the contact activation of the intrinsic blood coagulation pathway. APOH causes a reduction of the prothrombinase binding sites on platelets and reduces the activation caused by collagen when thrombin is present at physiological serum concentrations of APOH suggesting a regulatory role of APOH in coagulation. APOH plasma concentrations are strongly associated to metabolic syndrome alterations and vascular disease in type 2 diabetic and could be considered as a clinical marker of cardiovascular risk.

References

1. Schousboe I, 1985, Blood. 66 (5): 1086-91.
2. Nimpf J, et al., 1986, Biochim. Biophys. Acta 884 (1): 142-9.
3. Nimpf J, et al., 1987, Atherosclerosis. 63 (2-3): 109-14.
4. Ruii, G. et al., 1997, Clin Genet 52 (3): 167-72.
5. Castro, A. et al., 2009, Atherosclerosis. Oct 6.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>