

# Human REG4 / RELP / GISP Protein (Fc Tag)



**Sino Biological Inc.**  
Biological Solution Specialist

Catalog Number: 11186-H01H

## General Information

### Gene Name Synonym:

GISP; REG-IV; RELP; 2010002L15Rik; GISP; RELP

### Protein Construction:

A DNA sequence encoding the human REG4 (Q9BYZ8-1) (Asp23-Pro158) was expressed with the fused Fc region of human IgG1 at the N-terminus.

**Source:** Human

**Expression Host:** Human Cells

## QC Testing

**Purity:** > 90 % as determined by SDS-PAGE

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

**Predicted N terminal:** Glu

### Molecular Mass:

The recombinant human REG4/Fc is a disulfide-linked homodimer. The reduced monomer comprises 396 amino acids and has a predicted molecular mass of 44.3 kDa. The apparent molecular mass of the protein is approximately 45 kDa in SDS-PAGE under reducing conditions due to glycosylation.

### Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

Regenerating islet-derived protein 4, also known as REG-like protein, REG4, GISP and RELP, a member of the regenerating gene family belonging to the calcium (C-type) dependent lectin superfamily, has been found to be involved in malignancy in several different organs including the stomach, colorectum, pancreas and prostate. It is highly expressed in the gastrointestinal tract and markedly up-regulated in colon adenocarcinoma, pancreatic cancer, gastric adenocarcinoma, and inflammatory bowel disease. Expression of the Reg4 in different cell types has been associated with regeneration, cell growth and cell survival, cell adhesion and resistance to apoptosis. REG4 protein overexpression is associated with an unfavorable response to preoperative chemoradiotherapy and may be used as a predictive biomarker clinically. REG4 may play an important role in the development and progression of colorectal cancer, as well as in intestinal morphogenesis and epithelium restitution.

## References

- 1.Li FY, *et al.* (2010) RegIV expression showing specificity to gastrointestinal tract and its potential role in diagnosing digestive tract neuroendocrine tumor. *J Zhejiang Univ Sci B.* 11(4):258-66.
- 2.Rafa L, *et al.* (2010) REG4 acts as a mitogenic, motility and pro-invasive factor for colon cancer cells. *Int J Oncol.* 36(3): 689-98.
- 3.Hu G, *et al.* (2010) Purification of a bioactive recombinant human Reg IV expressed in *Escherichia coli*. *Protein Expr Purif.* 69(2): 186-90.

**Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.**

**Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>**