

Human TLK2 / PKU-ALPHA Protein (aa 397-772)

Catalog Number: 11551-HNCB



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

HsHPK; PKU-ALPHA; 4933403M19Rik; PKU-alpha; PKUalpha; Tlk

Protein Construction:

A DNA sequence encoding the human TLK2 (Q86UE8-1) (Leu397-Asn772) was fused with two additional amino acids (Gly & Pro) at the N-terminus.

Source: Human

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Kinase activity untested

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gly

Molecular Mass:

The recombinant human TLK2 consists of 378 amino acids and has a calculated molecular mass of 43.6 kDa. The recombinant protein migrates as an approximately 44 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Supplied as sterile 20mM Tris, 500mM NaCl, 3mM DTT, 10% glycerol, pH 8.0.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Serine / threonine-protein kinase tousled-like 2, also known as PKU-alpha, Tousled-like kinase 2 and TLK2, is a nucleus protein which belongs to the protein kinase superfamily and Ser/Thr protein kinase family. The tousled-like kinases are an evolutionarily conserved family of proteins implicated in DNA repair, DNA replication and mitosis in metazoans and plants. Their absence from the yeasts and other eukaryotic 'microbes' suggests a specific role for them in the development of multicellular organisms. Tousled-like kinase 2 / TLK2 is widely expressed. It is present in fetal placenta, liver, kidney, pancreas, heart and skeletal muscle. It is also found in adult cell lines. Tousled-like kinase 2 / TLK2 contains one protein kinase domain. Tousled-like kinase 2 / TLK2 is rapidly and transiently inhibited by phosphorylation following the generation of DNA double-stranded breaks during S-phase. This is cell cycle checkpoint and ATM-pathway dependent and appears to regulate processes involved in chromatin assembly.

References

- 1.Sillje HHW. et al., 2001,Curr Biol. 11: 1068-73.
- 2.Groth A. et al., 2003, EMBO J. 22: 1676-87.
- 3.Beausoleil SA. et al., 2004, Proc Natl Acad Sci. 101: 12130-5.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>