

Human PDE1C Protein (His & GST Tag)

Catalog Number: 13605-H20B



Sino Biological Inc.

Biological Solution Specialist

General Information

Gene Name Synonym:

cam-PDE1C; hCam-3; Hcam3

Protein Construction:

A DNA sequence encoding the human PDE1C (Q8TAE4) (Met1-Glu634) was fused with the N-terminal polyhistidine-tagged GST tag at the N-terminus.

Source: Human

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 85 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met

Molecular Mass:

The recombinant human PDE1C /GST chimera consists of 871 amino acids and has a calculated molecular mass of 100 kDa. The recombinant protein migrates as an approximately 97 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% gly, 3mM DTT

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

PDE1C belongs to the cyclic nucleotide phosphodiesterase family, PDE1 subfamily. Phosphodiesterases (PDEs) are a family of related phosphohydrolyases that selectively catalyze the hydrolysis of 3' cyclic phosphate bonds in adenosine and/or guanine 3',5' cyclic monophosphate (cAMP and/or cGMP). They regulate the cellular levels, localization and duration of action of these second messengers by controlling the rate of their degradation. PDEs are expressed ubiquitously, with each subtype having a specific tissue distribution. These enzymes are involved in many signal transduction pathways and their functions include vascular smooth muscle proliferation and contraction, cardiac contractility, platelet aggregation, hormone secretion, immune cell activation, and they are involved in learning and memory. PDE1C has a high affinity for both cAMP and cGMP. It is expressed in several tissues, including brain and heart. As a cyclic nucleotide phosphodiesterase, PDE1C has a dual-specificity for the second messengers cAMP and cGMP.

References

1. Strausberg RL, *et al.* (2003) Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. *Proc Natl Acad Sci.* 99(26):16899-903.
2. Dolci S, *et al.* (2006) Subcellular localization and regulation of type-1C and type-5 phosphodiesterases. *Biochem Biophys Res Commun.* 341(3):837-46.
3. Vandeput F, *et al.* (2007) Cyclic nucleotide phosphodiesterase PDE1C1 in human cardiac myocytes. *Biochemistry. J Biol Chem.* 282(45):32749-57.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>