

Human Cadherin-17 / LI-cadherin / CDH17 Protein (His Tag)

Catalog Number: 11360-H08H



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

CDH16; HPT-1; HPT1; HPT-1; HPT-1/LI

Protein Construction:

A DNA sequence encoding the human CDH17 (Q12864-1) extracellular domain (Met 1-Met 787) was expressed, fused with a polyhistidine tag at the C-terminus.

Source: Human

Expression Host: Human Cells

QC Testing

Purity: > 88 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gln 23

Molecular Mass:

The recombinant human CDH17 consists of 776 amino acids and predicts a molecular mass of 87 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rh CDH17 is approximately 105-115 kDa due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Cadherin-17 or LI-cadherin is a member of the cadherin superfamily, genes encoding calcium-dependent, membrane-associated glycoproteins. Cadherin-17/LI-cadherin is a cadherin-like protein consisting of an extracellular region, 7 cadherin domains, and a transmembrane region but lacking the conserved cytoplasmic domain. The protein is a component of the gastrointestinal tract and pancreatic ducts, acting as an intestinal proton-dependent peptide transporter in the first step in oral absorption of many medically important peptide-based drugs. The protein may also play a role in the morphological organization of liver and intestine. Alternative splicing of the encoding gene results in multiple transcript variants. Cadherin-17/LI-cadherin preferentially interact with themselves in a homophilic manner in connecting cells. Cadherin-17 may thus contribute to the sorting of heterogeneous cell types and have a role in the morphological organization of liver and intestine. It's also involved in intestinal peptide transport. Experiments have reported the association between Cadherin-17/LI-cadherin and gastric cancer. Cadherin-17/LI-cadherin expression was detected in 63/94 of gastric adenocarcinomas in addition to intestinal metaplasia. The expression of Cadherin-17 tended to be associated with intestinal type carcinoma, and carcinomas with Cadherin-17 expression was significantly more frequent in advanced stage cases than in early stage. Cadherin-17 is also a useful immunohistochemical marker for diagnosis of adenocarcinomas of the digestive system.

References

- 1.Liu LX, *et al.* (2009) Targeting cadherin-17 inactivates Wnt signaling and inhibits tumor growth in liver carcinoma. *Hepatology*. 50(5): 1453-63.
- 2.Ito R, *et al.* (2005) Clinicopathological significant and prognostic influence of cadherin-17 expression in gastric cancer. *Virchows Arch*. 447(4): 717-22.
- 3.Horsfield J, *et al.* (2002) Cadherin-17 is required to maintain pronephric duct integrity during zebrafish development. *Mech Dev*. 115(1-2): 15-26.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>