

Human IKB alpha / NFKBIA Protein (His Tag)



Sino Biological Inc.

Biological Solution Specialist

Catalog Number: 12045-H07E

General Information

Gene Name Synonym:

IKBA; MAD-3; NFKBI; AI462015; Nfkb1

Protein Construction:

A DNA sequence encoding the human NFKBIA (NP_065390.1) (Phe 2-Leu 317) was expressed, with a polyhistidine tag at the N-terminus.

Source: Human

Expression Host: E. coli

QC Testing

Purity: > 90 % as determined by SDS-PAGE

Endotoxin:

Please contact us for more information.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met

Molecular Mass:

The recombinant human NFKBIA consisting of 323 amino acids and has a calculated molecular mass of 36.4 kDa. It migrates as an approximately 38 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile 50mM Tris, 0.5M NaCl, pH 8.0

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha (I κ B alpha, NFKBIA, or IKBA), is a member of the NF-kappa-B inhibitor family that function to inhibit the NF-kB transcription factor. NFKBIA inhibits NF-kB by masking the nuclear localization signals (NLS) of NF-kB proteins and keeping them sequestered in an inactive state in the cytoplasm. In addition, NFKBIA blocks the ability of NF-kB transcription factors to bind to DNA, which is required for NF-kB's proper functioning. Signal-induced degradation of I kappa B alpha exposes the nuclear localization signal of NF-kappa B, thus allowing it to translocate into the nucleus and activate transcription from responsive genes. An autoregulatory loop is established when NF-kappa B induces expression of the I kappa B alpha gene and newly synthesized I kappa B alpha accumulates in the nucleus where it negatively regulates NF-kappa B-dependent transcription. As part of this post-induction repression, the nuclear export signal on I kappa B alpha mediates transport of NF-kappa B-I kappa B alpha complexes from the nucleus to the cytoplasm. Deletion of NFKBIA has an effect that is similar to the effect of EGFR amplification in the pathogenesis of glioblastoma and is associated with comparatively short survival. Polymorphisms in NFKBIA may be important in pre-disposition to and outcome after treatment, of multiple myeloma (MM). The NFKBIA gene product, IkappaBalp, binds to NF-kappaB preventing its activation and is important in mediating resistance to apoptosis in B-cell lymphoproliferative diseases.

References

- 1.Verma IM, *et al.* (1995) Rel/NF-kappa B/I kappa B family: intimate tales of association and dissociation. *Genes Dev.* 9 (22): 2723-35.
- 2.Jacobs MD, *et al.* (1998) Structure of an IkappaBalp/NF-kappaB complex. *Cell* 95 (6): 749-58.
- 3.Hay RT, *et al.* (1999) Control of NF-kappa B transcriptional activation by signal induced proteolysis of I kappa B alpha. *Philos Trans R Soc Lond B Biol Sci.* 354(1389): 1601-9.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>