

Human SerpinC1 / AntithrombinIII / ATIII Protein (His Tag)

Catalog Number: 10142-H08H



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

SerpinC1; AT3; AT3D; ATIII; MGC22579; SerpinC1; THPH7; A1114908; At-3; At3; ATIII

Protein Construction:

A DNA sequence encoding the human SerpinC1 (NP_000479.1) (Met 1-Lys 464) was expressed with a C-terminal polyhistidine tag.

Source: Human

Expression Host: Human Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to inhibit thrombin (Sigma, Catalog # T4648) cleavage of a fluorogenic peptide substrate Boc-VPR-AMC. The IC50 value is < 5 nM.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: His 33

Molecular Mass:

The recombinant human SerpinC1 consists of 443 amino acids and has a predicted molecular mass of 50.5 kDa. In SDS-PAGE, the apparent molecular mass of rhSerpinC1 is approximately 55-60 kDa due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

SerpinC1, also known as antithrombin III (AT III), is a member of the serpin superfamily of serine protease inhibitors, and has been found to be a marker for disseminated intravascular coagulation (DIC) and to be of prognostic significance in septic patients. SerpinC1 synthesized in the liver is the principal plasma serpin of blood coagulation proteases and inhibits thrombin and other factors such as Xa by the formation of covalently linked complexes. Thus it is one of the most important coagulation inhibitors and the fundamental enzyme for the therapeutic action of heparin. In common with SerpinA5 and D1, the inhibitory activity of SerpinC1 undergoes a dramatic increase in the presence of heparin and other glycosaminoglycans. ATIII mediates the promotion of prostaglandin release, an inhibitor of leucocyte activation and downregulator of many proinflammatory cytokines. Antithrombin III exerts anti-inflammatory properties in addition to its anti-coagulative mechanisms. In animal models of sepsis, ATIII affected cytokine plasma concentrations with a decrease of pro-inflammatory cytokines. The deficiency or functional abnormality of ATIII may result in an increased risk of thromboembolic disease, such as deep vein thrombosis and pulmonary embolism. In addition, it has been reported that SerpinC1 can alter or influence inflammatory processes via inhibition of NF-κB activation or actin polymerization.

References

1. de Sousa JC, *et al.* (1991) Antithrombin III. Physiologic, physiopathologic and laboratory aspects. *Rev Port Cardiol.* 10(9): 693-9.
2. Totzke G, *et al.* (2001) Antithrombin III enhances inducible nitric oxide synthase gene expression in vascular smooth muscle cells. *Cell Immunol.* 208(1): 1-8.
3. Ostermann H. (2002) Antithrombin III in Sepsis. New evidences and open questions. *Minerva Anesthesiol.* 68(5): 445-8.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>