

Mouse Neuroligin 1 / NLGN1 Protein (His Tag)



Sino Biological Inc.

Biological Solution Specialist

Catalog Number: 50725-M08H

General Information

Gene Name Synonym:

NL1; 6330415N05Rik; BB179718; mKIAA1070; NL1; Nlg1

Protein Construction:

A DNA sequence encoding the mouse NLGN1 isoform 1 (Q99K10-1) extracellular domain (Met 1-Ser 697) was expressed, with a C-terminal polyhistidine tag.

Source: Mouse

Expression Host: Human Cells

QC Testing

Purity: > 97 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gln 46

Molecular Mass:

The recombinant mouse NLGN1 consists of 671 amino acids and predicts a molecular mass of 73.5 KDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rmNLGN1 is approximately 100-110 KDa as a result of glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Neuroligin 1 (NLGN1) belongs to the type-B carboxylesterase/lipase family, is a synaptic cell-adhesion molecule that is enriched in postsynaptic densities where it may recruit receptors, channels, and signal-transduction molecules to synaptic sites of cell adhesion. Neuroligins consist of five members (NLGN1, NLGN2, NLGN3, NLGN4 and NLGN4Y), which interact with beta-neurexins and this interaction is involved in the formation of functional synapses. The extracellular domain of functional Neuroligin 1 associates as a dimer when analyzed by sedimentation equilibrium. Neuroligin 1 has a unique N-linked glycosylation pattern in the neuroligin family, and glycosylation and its processing modify neuroligin activity. Neuroligin 1 is a potent trigger for the de novo formation of synaptic connections, and it has recently been suggested that it is required for the maturation of functionally competent excitatory synapses. The persistent expression of Neuroligin 1 is required for the maintenance of NMDAR-mediated synaptic transmission, which enables normal development of synaptic plasticity and long-term memory in the amygdala of adult animals.

References

1. Song JY, *et al.* (1999) Neuroligin 1 is a postsynaptic cell-adhesion molecule of excitatory synapses. *Proc Natl Acad Sci U S A.* 96(3): 1100-5.
2. Comoletti D, *et al.* (2003) Characterization of the interaction of a recombinant soluble neuroligin-1 with neurexin-1beta. *J Biol Chem.* 278(50): 50497-505.
3. Ylisaukko-oja T, *et al.* (2005) Analysis of four neuroligin genes as candidates for autism. *Eur J Hum Genet.* 13(12): 1285-92.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>