

Mouse ENPEP / Aminopeptidase A Protein (His Tag)

Catalog Number: 50082-M07B



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

APA; CD249; ENPEP; gp160; 6030431M22Rik; APA; Bp-1/6C3; ENPEP; Ly-51; Ly51; ENPEP

Protein Construction:

A DNA sequence encoding the mature form of mouse ENPEP (P16406) (Arg41-Pro945) was expressed, with a polyhistidine tag at the N-terminus.

Source: Mouse

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 91 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to cleave the fluorogenic peptide substrate, Glu-7-amido-4-methyl coumarin. The specific activity is > 200 pmoles/min/μg.

Endotoxin:

< 1.0 EU per μg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: His

Molecular Mass:

The recombinant mouse ENPEP consists of 921 amino acids and predicts a molecular mass of 106 KDa. It migrates as an approximately 113 KDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

ENPEP, also known as aminopeptidase A, is a member of the peptidase M1 family. Members of this family are involved in response to cadmium ion and proteolysis. They located in 6 components and are expressed in 26 plant structures. ENPEP is expressed by epithelial cells of the proximal tubule cells and the glomerulus of the nephron. It also can be detected in a variety of other tissues. ENPEP probably plays a role in regulating growth and differentiation of early B-lineage cells. It also may play a role in the catabolic pathway of the renin-angiotensin system. ENPEP is a zinc-dependent membrane-bound aminopeptidase that catalyzes the cleavage of glutamatic and aspartatic amino acid residues from the N-terminus of polypeptides. It degrades vasoconstricting angiotensin II into angiotensin III and therefore helps to regulate blood pressure.

References

- 1.Speth RC, *et al.* (2008) The significance of brain aminopeptidases in the regulation of the actions of angiotensin peptides in the brain. *Heart Fail Rev.* 13(3):299-309.
- 2.Rose JE, *et al.* (2010) Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. *Mol Med.* 16(7-8):247-53.
- 3.Pérez I, *et al.* (2009) Increased APN/CD13 and acid aminopeptidase activities in head and neck squamous cell carcinoma. *Head Neck.* 31(10):1335-40.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>