

Mouse ICOS / AILIM / CD278 Protein (Fc Tag)



Sino Biological Inc.
Biological Solution Specialist

Catalog Number: 50466-M02H

General Information

Gene Name Synonym:

AILIM; CD278; CVID1; AILIM; CCLP; CRP-1; H4; Ly115

Protein Construction:

A DNA sequence encoding the mouse ICOS (Q9WVS0) extracellular domain (Met 1-Leu 142) was fused with the Fc region of human IgG1 at the C-terminus.

Source: Mouse

Expression Host: Human Cells

QC Testing

Purity: > 97 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Glu 21

Molecular Mass:

The recombinant mouse ICOS/Fc is a disulfide-linked homodimer. The reduced monomer consists of 363 amino acids and has a predicted molecular mass of 40.9 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of mICOS/Fc monomer is approximately 40-45 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Inducible costimulator (ICOS), also called AILIM (activation-inducible lymphocyte immunomediatory molecule) is a cell-surface receptor, and belongs to the CD28 family of immune costimulatory receptors consisting of CD28, CTLA-4 and PD-1. The interaction of B7-H2/ICOS plays a critical role in Th cell differentiation, T-B cell interactions which is essential for germinal center formation, and humoral immune responses, and as well as the production of cytokine IL-4. In addition, ICOS is more potent in the induction of IL-10 production, a cytokine important for suppressive function of T regulatory cells. The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway provides key second signals that can regulate the activation, inhibition and fine-tuning of T-lymphocyte responses. ICOS stimulates both Th1 and Th2 cytokine production but may have a preferential role in Th2 cell development. Moreover, The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway has been suggested of being involved in the development of airway inflammation and airway hyperresponsiveness.

References

- 1.Coyle AJ, *et al.* (2004) The role of ICOS and other costimulatory molecules in allergy and asthma. Springer Semin Immunopathol. 25(3-4): 349-59.
- 2.Chen YQ, *et al.* (2006) CD28/CTLA-4-CD80/CD86 and ICOS-B7RP-1 costimulatory pathway in bronchial asthma. Allergy. 61(1): 15-26.
- 3.van Berkel ME, *et al.* (2006) CD28 and ICOS: similar or separate costimulators of T cells Immunol Lett. 105(2): 115-22.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>