

Mouse DR6 / TNFRSF21 Protein (His Tag)

Catalog Number: 50263-M08H



Sino Biological Inc.

Biological Solution Specialist

General Information

Gene Name Synonym:

BM-018; CD358; DR6; AA959878; DR6; R74815; TR7

Protein Construction:

A DNA sequence encoding the mouse TNFRSF21 (NP_055267.1) extracellular domain (Met 1-His 349) was fused with a polyhistidine tag at the C-terminus.

Source: Mouse

Expression Host: Human Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Immobilized mouse TNFRSF21-His at 10 µg/ml (100 µl/well) can bind biotinylated human APP-Fc (Cat:10703-H02H with a linear range of 0.31-5 µg/ml).

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gln 42

Molecular Mass:

The secreted recombinant mouse TNFRSF21 comprises 319 amino acids and has a predicted molecular mass of 35 kDa. As a result of glycosylation, it migrates as an approximately 55-60 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

TNFRSF21 (death receptor-6, DR6) is an orphan TNF receptor superfamily member and belongs to a subgroup of receptors called death receptors. This type I transmembrane receptor possesses four extracellular cysteine-rich motifs and a cytoplasmic death domain. DR6 is an extensively posttranslationally modified transmembrane protein and that N- and O-glycosylations of amino acids in its extracellular part. DR6 interacts with the adaptor protein TRADD and mediates signal transduction through its death domain, and expression of DR6 in mammalian cells induces activation of both NF-kappaB and JNK and cell apoptosis. DR6 knockout mice have enhanced CD4+ T cell proliferation and Th2 cytokine production, suggested that DR6 serves as an important regulatory molecule in T-helper cell activation, and is involved in inflammation and immune regulation. DR6 is expressed ubiquitously with high expression in lymphoid organs, heart, brain and pancreas. Some tumor cells overexpress DR6, typically in conjunction with elevated anti-apoptosis molecules. DR6 may also be involved in tumor cell survival and immune evasion, which is subject to future investigations.

References

1. Pan G, *et al.* (1998) Identification and functional characterization of DR6, a novel death domain-containing TNF receptor. *FEBS Lett.* 431(3): 351-6.
2. Benschop R, *et al.* (2009) Tumor necrosis factor receptor superfamily member 21: TNFR-related death receptor-6, DR6. *Adv Exp Med Biol.* 647: 186-94.
3. Klma M, *et al.* (2009) Functional analysis of the posttranslational modifications of the death receptor 6. *Biochim Biophys Acta.* 1793(10): 1579-87.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>