

Mouse CD32 / FCGR2B Protein (His & AVI Tag), Biotinylated

Catalog Number: 50030-M27H-B



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

CD32; CD32A; CD32B; CDw32; Fc gamma RII; FCG2; FcGR; FCGR21; FCGR2A; IGFR2

Protein Construction:

A DNA sequence encoding the extracellular domain (Met 1-Arg 217) of mouse FGGR2B (NP_001070657.1) as fused with a c-terminal polyhistidine tagged AVI tag at the C-terminus. The expressed protein was biotinylated in vivo by the Biotin-Protein ligase (BirA enzyme) which is co-expressed.

Source: Mouse

Expression Host: Human Cells

QC Testing

Purity: > 97 % as determined by SDS-PAGE

Bio Activity:

1. Measured by its ability to bind human IgG1 in a functional ELISA. 2. Labeling ratio of biotin to protein: 0.5

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Thr 40

Molecular Mass:

The secreted recombinant mouse FGGR2B consists of 212 amino acids and has a predicted molecular mass of 23.9 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rm FGGR2B is approximately 38-43 kDa due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Receptors for Fc portion of IgG (Fcγ Rs) are members of the Ig superfamily, and are divided into three classes designated Fcγ RI (CD64), Fcγ RII (CD32), and Fcγ RIII (CD16). CD32 protein is a low affinity receptor for IgG that binds only IgG immune complexes and is expressed on a diverse range of cells such as monocytes, macrophages, neutrophils, eosinophils, platelets, and B cells. Human CD32 class is encoded by three closely related genes, and designated Fcγ RII A, B, and C which share 94-99% amino acid identity in their extracellular domains but differ substantially in their transmembrane and cytoplasmic domains. CD32 is involved in a number of immune responses including antibody-dependent cell-mediated cytotoxicity, clearance of immune complexes, release of inflammatory mediators, and regulation of antibody production.

References

1. Williams TE, *et al.* (2000) Concurrent and independent binding of Fcγ receptors IIa and IIb to surface-bound IgG. *Biophys J.* 79(4): 1867-75.
2. Kanfers D, *et al.* (2007) Expression of activated Fc gamma RII discriminates between multiple granulocyte-priming phenotypes in peripheral blood of allergic asthmatic subjects. *J Allergy Clin Immunol.* 120(5): 1073-81.
3. Veri MC, *et al.* (2007) Monoclonal antibodies capable of discriminating the human inhibitory Fcγ receptor IIB (CD32B) from the activating Fcγ receptor IIA (CD32A): biochemical, biological and functional characterization. *Immunology.* 121(3): 392-404.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>