

Rat JAM-A / F11R Protein (Fc Tag)

Catalog Number: 80240-R02H



Sino Biological Inc.

Biological Solution Specialist

General Information

Gene Name Synonym:

CD321; JAM; JAM1; JAMA; JCAM; KAT; PAM-1; 9130004G24; AA638916; ESTM33; JAM; JAM-1; JAM-A; Jcam; Jcam1; Ly106

Protein Construction:

A DNA sequence encoding the rat F11R () was expressed, fused with the Fc region of human IgG1 at the C-terminus.

Source: Rat

Expression Host: Human Cells

QC Testing

Purity: > 96 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Lys 27

Molecular Mass:

The recombinant rat F11R/Fc is a disulfide-linked homodimer. The reduced monomer comprises 453 amino acids and has a predicted molecular mass of 49.9 kDa. The apparent molecular mass of the protein is approximately 60 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Junctional adhesion molecule-A (JAM-A), also known as F11 receptor (F11R) or Cluster of Differentiation 321 (CD321), is a transmembrane protein expressed at tight junctions of epithelial and endothelial cells, as well as on circulating leukocytes. JAM-A protein serves as a serotype-independent receptor for mammalian orthoreoviruses (reoviruses). It is also a ligand for the integrin LFA1, involves in leukocyte transmigration. As a cell adhesion molecule of the immunoglobulin superfamily, JAM-A protein involves in platelet adhesion, secretion and aggregation, and plays a crucial role in inflammatory thrombosis and atherosclerosis. In addition, it may be a potential therapeutic target for breast cancer.

References

1. Guglielmi KM, *et al.* (2007) Reovirus binding determinants in junctional adhesion molecule-A. *J Biol Chem.* 282(24): 17930-40.
2. Yeung D, *et al.* (2008) Decreased junctional adhesion molecule-A expression during blood-brain barrier breakdown. *Acta Neuropathol.* 115(6): 635-42.
3. Ong KL, *et al.* (2009) Elevated plasma level of soluble F11 receptor/junctional adhesion molecule-A (F11R/JAM-A) in hypertension. *Am J Hypertens.* 22(5): 500-5.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>