

Rat ALCAM / CD166 Protein (Fc Tag)

Catalog Number: 80221-R02H



Sino Biological Inc.

Biological Solution Specialist

General Information

Gene Name Synonym:

CD166; MEMD; AI853494; BEN; CD166; DM-GRASP; MuSC; SC1

Protein Construction:

A DNA sequence encoding the rat ALCAM (O35112) (Met1-Lys527) was expressed with the Fc region of human IgG1 at the C-terminus.

Source: Rat

Expression Host: Human Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Immobilized rat CD6-His (Cat:80311-R08H) at 10 µg/ml (100 µl/well) can bind rat ALCAM-Fc3, The EC₅₀ of rat ALCAM-Fc3 is 19.5-45.5 ng/ml.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Trp 28

Molecular Mass:

The recombinant rat ALCAM/Fc is a disulfide-linked homodimer. The reduced monomer comprises 741 amino acids and has a predicted molecular mass of 83.1 kDa. The apparent molecular mass of the protein is approximately 94 kDa in SDS-PAGE under reducing conditions due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Activated leukocyte cell adhesion molecule (ALCAM)/Cluster of differentiation (CD166) is a type I transmembrane cell adhesion molecule belonging to the Ig superfamily and a ligand for CD6 that is expressed on T lymphocytes. The extracellular domain of ALCAM contains five Ig-like domains (three Ig-like C2-type domains and two Ig-like V-type domains), of which the amino-terminal V1 domain is essential for ligand binding and ALCAM-mediated cell aggregation. ALCAM mediates both heterophilic (ALCAM-CD6) and homophilic (ALCAM-ALCAM) cell-cell interactions. ALCAM/CD6 interaction plays a role in T cell development and T cell regulation, as well as in the binding of T- and B-cells to activated leukocytes. Recently, homophilic (ALCAM-ALCAM) adhesion was shown to play important roles in tight cell-to-cell interaction and regulation of stem cell differentiation. While expressed in a wide variety of tissues, ALCAM is usually restricted to subsets of cells involved in dynamic growth and/or migration, including neural development, branching organ development, hematopoiesis, immune response and tumor progression. And CD166 is regarded as a potential novel breast cancer indicator and therapeutic target.

References

1. Swart GW. (2002) Activated leukocyte cell adhesion molecule (CD166/ALCAM): developmental and mechanistic aspects of cell clustering and cell migration. *Eur J Cell Biol.* 81(6): 313-21.
2. Fujiwara H, *et al.* (2003) Human blastocysts and endometrial epithelial cells express activated leukocyte cell adhesion molecule (ALCAM/CD166). *J Clin Endocrinol Metab.* 88(7): 3437-43.
3. Jezierska A, *et al.* (2006) ALCAM/CD166 protects breast cancer cells against apoptosis and autophagy. *Med Sci Monit.* 12(8): BR263-73.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>