

Cynomolgus BTC / Betacellulin Protein (Fc Tag)

Catalog Number: 90033-C02H



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

Bcn

Protein Construction:

A DNA sequence encoding the cynomolgus BTC (XP_001101880.1) (Met1-Tyr111) was expressed with the Fc region of human IgG1 at the C-terminus.

Source: Cynomolgus

Expression Host: Human Cells

QC Testing

Purity: > 85 % as determined by SDS-PAGE

Bio Activity:

Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED₅₀ for this effect is typically 5-40 ng/mL.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Asp 32

Molecular Mass:

The recombinant cynomolgus BTC is a disulfide-linked homodimer. The reduced monomer comprises 321 amino acids and has a calculated molecular mass of 36 KDa. The apparent molecular mass of the protein is approximately 46 KDa respectively in SDS-PAGE.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Betacellulin(BTC) is a member of the epidermal growth factor (EGF) family. These soluble proteins are ligands for one or more of the four receptor tyrosine kinases encoded by the ErbB gene family (ErbB-1/epidermal growth factor receptor (EGFR), neu/ErbB-2/HER2, ErbB-3/HER3 and ErbB-4/HER4). Betacellulin is a 32-kilodalton glycoprotein that appears to be processed from a larger transmembrane precursor by proteolytic cleavage. This protein is a ligand for the EGF receptor. BTC is a polymer of about 62-111 amino acid residues. Secondary Structure: 6% helical (1 helices; 3 residues)36% beta sheet (5 strands; 18 residues). BTC was originally identified as a growth-promoting factor in mouse pancreatic β-cell carcinoma cell line and has since been identified in humans. It plays a role in the growth and development of the neonate and/or mammary gland function. Betacellulin is a potent mitogen for retinal pigment epithelial cells and vascular smooth muscle cells.

References

1. Shing Y, *et al.* (1993) Betacellulin: a mitogen from pancreatic beta cell tumors. *Science* . 259(5101): 1604-7.
2. Riese DJ, *et al.* (1996) Betacellulin activates the epidermal growth factor receptor and erbB-4, and induces cellular response patterns distinct from those stimulated by epidermal growth factor or neuregulin-beta. *Oncogene*. 12(2): 345-53.
3. Bastian SE, *et al.* (2001) Measurement of betacellulin levels in bovine serum, colostrum and milk. *J Endocrinol* . 168: 203-12.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>