

Cynomolgus / Rhesus M-CSF / CSF-1 Protein (Fc Tag)

Catalog Number: 90029-C02H



Sino Biological Inc.
Biological Solution Specialist

General Information

Gene Name Synonym:

CSF-1; M-CSF; MCSF; C87615; Csfm; MCSF; op

Protein Construction:

A DNA sequence encoding the cynomolgus CSF1 (XP_005542489.1) (Met1-Asn190) was expressed with the Fc region of human IgG1 at the C-terminus. Cynomolgus and Rhesus CSF1 sequences are identical.

Source: Cynomolgus

Expression Host: Human Cells

QC Testing

Purity: > (15.1+79.5+4.6) % as determined by SDS-PAGE.

Endotoxin:

< 1.0 EU per µg protein as determined by the LAL method.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Glu 33

Molecular Mass:

The recombinant cynomolgus CSF1 consists of 396 amino acids and predicts a molecular mass of 45.1 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Macrophage colony-stimulating factor 1, also known as CSF-1, M-CSF, Lanimostim and CSF1, is a single-pass membrane protein which is disulfide-linked as a homodimer or heterodimer. Granulocyte / macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. M-CSF/CSF-1 is known to facilitate monocyte survival, monocyte-to-macrophage conversion, and macrophage proliferation. M-CSF/CSF-1 is a secreted cytokine which influences hemopoietic stem cells to differentiate into macrophages or other related cell types. It binds to the Colony stimulating factor 1 receptor. M-CSF/CSF-1 may also be involved in development of the placenta. The active form of M-CSF/CSF-1 is found extracellularly as a disulfide-linked homodimer, and is thought to be produced by proteolytic cleavage of membrane-bound precursors. M-CSF/CSF-1 induces cells of the monocyte/macrophage lineage. It also plays a role in immunological defenses, bone metabolism, lipoproteins clearance, fertility and pregnancy. Upregulation of M-CSF/CSF-1 in the infarcted myocardium may have an active role in healing not only through its effects on cells of monocyte/macrophage lineage, but also by regulating endothelial cell chemokine expression.

References

1. Pandit J. et al., 1992, Science. 258: 1358-62.
2. Tokai M. et al., 2000, J Bacteriol. 182 (10): 2865-8.
3. Fan X. et al., 2001, Am J Physiol Endocrinol Metab. 280 (1): E103-11.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>