

# Rhesus CD16 / FCGR3 Protein (His Tag)

Catalog Number: 90013-C08H



**Sino Biological Inc.**  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

FCGR3

### Protein Construction:

A DNA sequence encoding the rhesus CD16 (NP\_001258584.1) extracellular domain (Met 1-Gly 206) was fused with a polyhistidine tag at the C-terminus.

**Source:** Rhesus

**Expression Host:** Human Cells

## QC Testing

**Purity:** > 95 % as determined by SDS-PAGE

### Bio Activity:

Measured by its binding ability in a functional ELISA. Immobilized Rhesus FCGR3-His (cat:90013-C08H) at 10 µg/ml (100 µl/well) can bind recombinant human IgG1 (Fc) (Cat:10702-HNAC). The  $EC_{50}$  of human IgG1 (Fc) (Cat:10702-HNAC) is 0.35-0.81 µg/ml.

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

**Predicted N terminal:** Gly 17

### Molecular Mass:

The recombinant rhesus CD16 consists of 201 amino acids and has a calculated molecular mass of 22.7 kDa.

### Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

Fc receptors bind the most common class of antibody, IgG, are called Fc gamma receptors (FcγR). FcγR is divided into three classes, Fc γ RI (CD64), Fc γ RII (CD32), and Fc γ RIII (CD16). CD16 protein is a multifunctional, low/intermediate affinity receptor, which belongs to the immunoglobulin superfamily. It is found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. Mouse CD16 is encoded by a single gene, while, human CD16 is expressed as two distinct forms (CD16a/FcγRIIIa and CD16b/FcγRIIIb) encoded by two different highly homologous genes in a cell type-specific manner. CD16 is involved in phagocytosis, secretion of enzymes, inflammatory mediators, antibody-dependent cellular cytotoxicity (ADCC), and clearance of immune complexes.

## References

1. Edberg JC, *et al.* (2002) Genetic linkage and association of Fc gamma receptor IIIA (CD16A) on chromosome 1q23 with human systemic lupus erythematosus. *Arthritis Rheum.* 46(8): 2132-40.
2. Li P, *et al.* (2002) Recombinant CD16A-Ig forms a homodimer and cross-blocks the ligand binding functions of neutrophil and monocyte Fc gamma receptors. *Mol Immunol.* 38(7): 527-38.
3. Li P, *et al.* (2007) Affinity and kinetic analysis of Fc gamma receptor IIIa (CD16a) binding to IgG ligands. *J Biol Chem.* 282(9): 6210-21.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>