

Canine CTLA-4 / CD152 Protein (ECD, Fc Tag)



Sino Biological Inc.
Biological Solution Specialist

Catalog Number: 70113-D02H

General Information

Gene Name Synonym:

ALPS5; CD; CD152; CELIAC3; CTLA-4; GRD4; GSE; IDDM12; Cd152; CtlA-4; Ly-56

Protein Construction:

A DNA sequence encoding the canine CD152 (NP_001003106.1) (Met1-Asp161) was expressed with the Fc region of human IgG1 at the C-terminus.

Source: Canine

Expression Host: Human Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE.

Endotoxin:

< 1.0 EU per µg protein as determined by the LAL method.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Lys 36

Molecular Mass:

The recombinant canine CD152 consists 364 amino acids and predicts a molecular mass of 40.1 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Cytotoxic T-lymphocyte protein 4, also known as CTLA4 and CD152, is a single-pass type I membrane protein and a member of the immunoglobulin superfamily. It is the second member of the CD28 receptor family. The ligands or counterreceptors for these two proteins are the B7 family members, CD80 (B7-1) and CD86 (B7-2). CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T cells and may play an important role in their functions. CD152 or cytotoxic T lymphocyte antigen-4 (CTLA-4) is an essential receptor involved in the negative regulation of T cell activation. Because of its profound inhibitory role, CD152 has been considered a sound susceptible candidate in autoimmunity and a persuasive target for cancer immunotherapy. In particular, recent evidence suggests that CD152 is also important in the homeostasis and function of a population of suppressive cells, termed regulatory T cells (Treg).

References

- 1.Slavik JM, *et al.* (1999) CD28/CTLA-4 and CD80/CD86 families: signaling and function. *Immunol Res.* 19(1): 1-24.
- 2.Holmberg D, *et al.* (2005) CTLA-4 (CD152) and its involvement in autoimmune disease. *Autoimmunity.* 38(3): 225-33.
- 3.Chin LT, *et al.* (2008) Immune intervention with monoclonal antibodies targeting CD152 (CTLA-4) for autoimmune and malignant diseases. *Chang Gung Med J.* 31(1): 1-15.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

Fax :+86-10-51029969 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>